

## Drug Facts

<b>Active ingredients</b>	<b>Purpose</b>
---------------------------	----------------

Naphazoline hydrochloride (0.012%).....	Redness reliever
Polyethylene glycol 300 (0.2%).....	Lubricant

### Uses

- relieves redness of the eye due to minor eye irritations
- temporary relief of burning and irritation due to dryness of the eye
- protects against further irritation

### Warnings

**For external use only**

#### Do not use

- if solution changes color or becomes cloudy

#### Ask a doctor before use if you have

- narrow angle glaucoma

#### When using this product

- do not touch tip of container to any



## Drug Facts (continued)

- surface to avoid contamination
- do not overuse as it may produce increased redness of the eye
- pupils may become enlarged temporarily
- remove contact lenses before using
- replace cap after use

#### Stop use and ask a doctor if

- you experience eye pain, changes in vision, continued redness or irritation of the eye
- condition worsens or persists for more than 72 hours

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

- instill 1 to 2 drops in the affected eye(s) up to four times daily

### Other information

- store at 15°-25°C (59°-77°F)
- keep tightly closed
- use before expiration date marked on the carton and bottle

### Inactive ingredients

benzalkonium chloride (0.01%), boric acid, edetate disodium, sodium borate, sodium chloride. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.



## Drug Facts (continued)

### Questions or comments?

( Call: 1-800-553-5340