

## ***Drug Facts***

| <b><i>Active ingredient</i></b>            | <b><i>Purpose</i></b> |
|--|-----------------------|
| Sodium chloride 5%.....Hypertonicity agent |                       |

### ***Uses***

temporary relief of corneal edema

### ***Warnings***

**For external use only**

#### **Do not use**

- except under the advice and supervision of a doctor

#### **When using this product**

- it may cause temporary burning and irritation
- replace cap after use
- to avoid contamination, do not touch tip of container to any surface

#### **Stop use and ask a doctor if**

- condition worsens or persists for more than 72 hours
- you experience eye pain, changes in vision, continued redness or irritation of the eye

#### **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

### ***Directions***

- pull down the lower lid of the affected eye
- apply a small amount (1/4 inch) of ointment to the inside of eyelid
- apply every 3 or 4 hours or as directed by a doctor ▶

## ***Drug Facts***

(continued)

### ***Other information***

- store at 15–25°C (59–77°F)
- keep tightly closed
- **DO NOT FREEZE**
- see crimp of tube or carton for Lot Number and Expiration Date
- do not use if difficult to dispense or visible particles are seen in the product
- serious side effects associated with use of the product may be reported to the phone number provided below

### ***Inactive ingredients***

lanolin, mineral oil, purified water, white petrolatum.

### ***Questions or comments?***

( Call 1-800-553-5340